ADVERSE CHILDHOOD EXPERIENCES (ACEs)

Daniel M. Christy, MPA, CPRP, Director WV Health Statistics Center December 15, 2015











What are ACEs?



Adverse

Childhood

Experience

An adverse childhood experience, or ACE, describes a **traumatic event** in a person's life occurring before the age of 18 that the person recalls as an adult.

What are ACEs? (continued)





Adverse Childhood Experiences

ACEs are experiences in childhood that are unhappy, unpleasant and/or hurtful.



ACEs are sometimes referred to as toxic stress or childhood trauma.

ACE Definition



An ACE is defined as surviving any of the following categories of abuse, neglect, or loss prior to age 18:

- Emotional abuse by a parent
- Physical abuse by a parent
- Sexual abuse by anyone
- Emotional neglect
- Physical neglect
- Loss of a parent
- Growing up with an alcohol and/or drug abuser in the household
- Living with a family member experiencing mental illness
- Experiencing the incarceration of a household member

ACE Study



Several ACE studies reveal strong relationships between ACEs and many later-life health risks and conditions. For example, higher ACE scores are associated with obesity, substance abuse, depression and lung, heart and liver disease. ACEs are also linked to homelessness and criminal justice involvement.

Integral Theory explains how ACEs can derail developmental processes as well as how community resources that foster resilience can help restore healthy development.

Health Problems Associated with ACE



According to the Centers for Disease Control and Prevention (CDC, 2014), ACEs are associated with the following risk factors, chronic disease, and health outcomes:

- Alcoholism and alcohol abuse
- Chronic obstructive pulmonary disease
- Depression
- Fetal death
- Health-related quality of life
- Illicit drug use
- Ischemic heart disease
- Risk for intimate partner violence

- Liver disease
- Multiple sexual partners
- Sexually transmitted diseases
- Smoking
- Suicide attempts
- Unintended pregnancies
- Early initiation of smoking
- Early initiation of sexual activity
- Adolescent pregnancy

ACE Categories



There is a total of 11 questions on the ACE module of the Behavioral Risk Factor Surveillance System (BRFSS) questionnaire. The 11 questions were combined to create 8 categories. The household alcohol and drug use questions were combined to create a household substance abuse category. The 3 questions on sexual abuse and rape were combined to create a sexual abuse category. The final 8 categories are:


- Substance Abuse
- Separation/Divorce
- Verbal Abuse
- Mental Illness

- Domestic Violence
- Physical Abuse
- Sexual Abuse
- Incarceration

Types of ACEs Among West Virginia Adults




Prevalence of Adverse Childhood Experiences Among West Virginia Adults, 2014



Number of ACEs Among West Virginia Adults




Number of Adverse Childhood Experiences Reported Among West Virginia Adults, 2014



High Risk by Gender




Prevalence of High Risk Among West Virginia Adults by Gender, 2014



High Risk by Age




Prevalence of High Risk Among West Virginia Adults by Age Group, 2014



High Risk by Education




Prevalence of High Risk Among West Virginia Adults by Education, 2014



High Risk by Income




Prevalence of High Risk Among West Virginia Adults by Income, 2014



Health Risk Behavior




Comparison of Health Risk Behavior Among Those at High Risk from ACEs and Those at Low Risk from ACEs



Chronic Diseases



Comparison of Chronic Diseases Among Those at High Risk from ACEs and Those at Low Risk from ACEs



Conclusions



West Virginia adults reported an average of 1.2 ACEs during their childhood and more than half report zero ACEs. In West Virginia, the most common ACE was household substance abuse, followed by separation/divorce, verbal abuse, mental illness, domestic violence, physical abuse, sexual abuse, and incarceration of a household member.

Approximately 17.6% or 228,739 West Virginia adults would be considered high risk for having or developing health problems based on having three or more ACEs.

Conclusions (continued)



West Virginia's statistics are comparable to the CDC's findings in that multiple ACEs are associated with fair/poor health, current smoking, binge drinking, obesity, chronic obstructive pulmonary disease, depression, and disability.

These findings lay a foundation for prevention of chronic disease and unhealthy behaviors based on modification or avoidance of ACEs.

What Can Reduce ACE Impacts?



What types of community and family skills reduce ACEs?

- Nurturing and positive relationships
- Knowledge of parenting and child development
- Parental resilience
- Social connections
- Concrete support in time of need

More Information and Resources



National Center for Trauma-Informed Care www.mentalhealth.samhsa.gov/nctic

National Child Traumatic Stress Network
www.nctsnet.org

Center for the Study of Social Policy - Information on Strengthening Families and Protective Factors

www.cssp.org

Center for Injury Prevention and Control www.cdc.gov/violenceprevention

Contact



West Virginia Department of Health and Human Resources
Bureau for Public Health
Health Statistics Center
350 Capitol Street, Room 165
Charleston, WV 25301
www.wvdhhr.ogr/bph/hsc

Daniel Christy, Director daniel.m.christy@wv.wv.gov (304) 356-4186

Tonya Yablonsky, Epidemiologist <u>Tonya.a.yablonsky@wv.gov</u> (304) 356-4187